

# MCQS

**1- Juvenile Hyaline Fibromatosis (J.H.F) is characterized by - mark the statement True (T) or False (F).**

- 1- Autosomal dominant inheritance  
T F
- 2- Large cutaneous nodules around head and neck gingival hyperplasia, joint contracture  
T F
- 3- The collagen present is type IV  
T F
- 4- The collagen present is mainly the alpha three VI  
T F
- 5- The main glycosaminoglycan present in tissue is Hyaluronan  
T F

**2- The differential diagnosis of J.H.F. includes :**

- 1- Lipoid proteinosis
- 2- Farber lipogranulomatosis
- 3- Mucopolysaccharidosis
- 4- Congenital generalized fibromatosis
- 5- Winchester syndrome

**3- Treatment tried for J.H.F. include (mark the correct answer) :**

- 1- Penicillin
- 2- Proteolytic enzymes
- 3- Ketotifen
- 4- Calcitriol
- 5- Dimethyl sulfoxide

**Give single best response (4 to 18) :**

**4 - Nonvenereal treponematoses in comparison with venereal syphilis**

- a. are caused by distinct organisms
- b. do not have a relapsing course
- c. are best treated with aminoglycosides
- d. tend to predominantly affect children
- e. are generally less destructive

**5- Yaws is caused by Treponema pallidum subsp.**

- a. endemicum
- b. pertenue
- c. pallidum
- d. carateum
- e. yertenu

**6- Secondary yaws lesions( daughter ) yaws**

- a. are not infectious
- b. are characteristically destructive
- c. are frequently located adjacent to body orifices
- d. tend to be dry and scant
- e. spare the palm and soles

**7- Endemic syphilis ( bejel )**

- a. is prevalent in dry arid climates
- b. was eradicated from Africa in the 1963
- c. affects males more than females
- d. tends to affect children 2 months to 2 years of age
- e. is transmitted through breast milk

**8- Characteristic clinical features of endemic syphilis include the following, except**

- a. mucous patches most common initial manifestation
- b. bone lesions spare extremities
- c. primary lesions encountered rarely
- d. destructive lesions of the palate and septum
- e. sparing of the cardiac system

**9- Pinta is unique among the treponematoses as it**

- a. has only cutaneous manifestations
- b. is found only on the Eastern hemisphere
- c. was eradicated in 1950s
- d. have no treatment
- e. affect mainly the elderly.



**10- The drug of choice for the nonvenereal treponematoses is**

- a. erythromycin
- b. penicillin G procaine
- c. chloramphenicol
- d. penicillin G benzathine
- e. tetracycline

**Select the one lettered item most closely related to each numbered item**

- a. Yaws
- b. Endemic syphilis
- c. Pinta
- d. All of the above
- e. None of the above

**11- positive rapid plasma reagin**

**12- chloramphenicol drug of choice**

**13- Does not affect eyes and central nervous system**

**14- Easily distinguished from venereal syphilis**

**15- Mucous patches**

**Choose the lettered item for each numbered item**

- a. Yaws
- b. Pinta
- c. Both
- d. Neither

**16- Inflammatory infiltrate composed mainly of plasma cells and lymphocytes**

**17- Acanthotic epidermis**

**18- Absence of melanin in the basal layer**

9- a.	11- d.	13- c.	15- b.	17- c.
10- d.	12- c.	14- c.	16- c.	18- b.
8- b.	10- d.	12- c.	14- c.	16- c.
2- T	4- T	5- F	1, 2, 3, 4, 5	3- 2, 3, 4, 5
1- F	3- F	4- 1, 2, 3, 4, 5	4- d.	6- c.
ANSWERS TO MCQS :	2- T	7- a.	5- b.	